## Medication interactions with warfarin (Coumadin®)

Many drugs interfere with warfarin. Make sure you notify all of your health care providers that you are taking warfarin so they will know which medications can be prescribed for you. It is also important to notify your VA Coumadin Clinic with any changes in your medications or prior to starting any new medications.

## How should I store my warfarin (Coumadin®)

Keep your warfarin in the container it came in, tightly closed, and out of reach of children. Store it at room temperature and away from light.

#### **VA Coumadin Clinics**

Albany (518) 626-5000

Bath (607) 664-4465

Canandaigua (585) 393-7137

Rochester (585) 463-2699

Syracuse (315) 425-2615

VA Western NY Healthcare System

Buffalo Pharmacist (716) 862-7212

Patients A-L (716) 862-8764

Patients M-Z (716) 862-8763

Batavia (585) 297-1110

#### Over the counter medications

### You <u>may</u> take these nonprescription medications with warfarin (Coumadin®):

acetaminophen (Tylenol®)

Chlor-Trimeton®

Contac®

Drixoril®

\*Maalox®

\*Mylanta®

Robitussin® Cough Syrup

Sudafed®

Tylenol Cold and Sinus®

# You may **not** take these nonprescription medications with warfarin (Coumadin®) without your health care provider's approval:

Alka-Seltzer®

aspirin (Bayer®, Bufferin®, Ecotrin®)

 $Excedrin^{\scriptsize{\circledR}}$ 

ibuprofen (Advil®, Motrin®)

Naproxen (Naprosyn®, Aleve®)

Nyquil Cold®

Pepto-Bismol @

### Potential interactions between herbs and warfarin (Coumadin®)

Angelica Root Goldenseal
Arnica Flower Green Tea

Anise Horse Chestnut

Asafoetida Licorice Root
Bogbean Lovage Root
Borage Seed Oil Meadowsweet

Bromelain Onion

Capsicum Papain
Celery Parsley

Chamomile Passionflower

Clove Poplar

Co-enzyme Q<sub>10</sub> Quassia

Cranberry Juice Red Clover

Dandelion Rue

Danshen St. John's Wart

Devil's Claw Sweet Clover

Dong Quai Sweet Woodruff

Fenugreek Tonka Beans

Feverfew Turmeric

Garlic Vitamin E

Ginger Willow Bark

Ginko Yarrow

Ginseng







<sup>\*</sup> **Note:** If you use antacids, such as Maalox® or Mylanta®, take them at least two hours before or two hours after taking your warfarin.

# Common prescription drug interactions with warfarin (Coumadin®)

Alcohol

Antianxiety drugs

Antiarrhythmics amiodarone

(Cordarone®)

Anticonvulsants

phenytoin (Dilantin®)

carbamazepine (Tegretol®)

Antibiotics

penicillin

ciprofloxacin

trimethoprim/

sulfamethoxazole (Bactrim®)

**Antidepressants** 

Antigout drugs

allopurinol (Zvloprim®)

Antineoplastics cancer drugs

Antiplatelet drugs

clopidogrel (Plavix®) ticlopidine (Ticlid®)

#### Antithyroid drugs

levothyroxine

(Levothroid $^{\mathbb{R}}$ ,

Synthroid®)

#### Beta-Adrenergic

**Blockers** 

propranolol (Inderal®)

Diabetes drugs (Oral)

#### **Diuretics**

furosemide (Lasix®)

#### Gastrointestinal

cimetidine (Tagamet®) omeprazole

(Prilosec®)

#### **Hypolipidemics**

simvastatin (Zocor®) atorvastatin (Lipitor®)

#### **Narcotics**


hydrocodone

(Lortab®, Vicodin®)

#### **Steroids**

prednisone

### **Reaching Us Is Easy**



#### Veterans Service Contact Center 1-888-823-9656

For information on eligibility, VA health care benefits, enrollment, or questions on your billing statement

#### Telcare 1-888-838-7890

For immediate medical advice any time any place

#### **VA Health Care on the Web**

www.va.gov/visns/visn02

For reliable health information on the Web

www.myhealth.va.gov

Produced by VA Healthcare Network Upstate New York / Public Affairs Office
Designed and printed exclusively by the Network 2 Print Shop, Bath VA Medical Center.
All rights reserved.

8/06









What you need to know about

Drug interactions

with warfarin

(Coumadin®)



